


CP-673451

Chemical Properties

CAS No. : 343787-29-1
 Formula: C₂₄H₂₇N₅O₂
 Molecular Weight: 417.5
 Appearance: no data available
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	CP-673451 is a specific inhibitor of PDGFRα/β (IC ₅₀ : 10/1 nM) with antiangiogenic and antitumor activity and the selectivity is higher 450-fold than other angiogenic receptors.
Targets(IC ₅₀)	VEGFR,PDGFR,c-Kit
In vitro	CP 673451 is a selective inhibitor of PDGFRα/β with IC ₅₀ of 10 nM/1 nM, exhibits >450-fold selectivity over other angiogenic receptors. In glioblastoma tumors, CP-673451 (33 mg/kg) provides >50% inhibition of PDGFR-β receptor for 4 hours corresponding to an EC ₅₀ of 120 ng/mL in plasma at C _{max} . In a sponge angiogenesis model, CP-673451 inhibits 70% of PDGF-BB-stimulated angiogenesis at a dose of 3 mg/kg (q.d. ×5, p.o., corresponding to 5.5 ng/mL at C _{max}). [1] CP-673451 decreases cell proliferation rate through mechanisms involving reduced phosphorylation of GSK-3α and GSK-3β. In both RD and RUCH2 cultures, CP-673451 impairs rhabdosphere-forming capacity and cell differentiation, causes increased senescence. [2]
In vivo	CP 673451 (once-daily p.o.) inhibits tumor growth (ED ₅₀ < 33 mg/kg) in a number of human tumor xenografts grown s.c. in athymic mice, including H460 human lung carcinoma, Colo205 and LS174T human colon carcinomas, and U87 mg human glioblastoma multiforme. [1] In RUCH2 xenograft-bearing mice, CP 673451 reduces tumor growth and stromal cell infiltration. [2]
Kinase Assay	Kinase inhibition assay: A glutathione S-transferase-tagged kinase domain construct of the intracellular portion of the PDGFR-β (amino acids 693-1401, accession no. J03278) is expressed in Sf-9 cells (baculovirus expression system). Enzyme kinetics are determined by incubating the enzyme with increasing concentrations of ATP in phosphorylation buffer [50 mmol/L HEPES (pH 7.3), 125 mmol/L NaCl, 24 mmol/L MgCl ₂ in Nunc Immuno MaxiSorp 96-well plates previously coated with 100 μL of 100 μg/mL poly-Glu-Tyr (4:1 ratio) diluted in PBS. After 10 minutes, the plates are washed (PBS, 0.1% Tween 20), incubated with anti-phosphotyrosine-horseradish peroxidase antibody, and diluted in PBS, 0.05% Tween 20, 3% BSA for 30 minutes at room temperature. The plates are washed as above and incubated with 3,3',5,5'-tetramethylbenzidine. The reaction is stopped by adding an equal volume of 0.09 NaH ₂ SO ₄ . The phosphotyrosine-dependent signal is then quantitated on a plate reader at 450 nm. For routine enzyme assays, the enzyme is incubated with 10 μM (final) ATP in the presence of compound diluted in DMSO (1.6% v/v DMSO assay final) for 30 minutes at room temperature in plates, as above, previously coated with 100 μL of 6.25 μg/mL poly-Glu-Tyr. The remainder of the assay is carried out as above, and IC ₅₀ values are calculated as percent inhibition of

	control.
Cell Research	PAE cells stably expressing full-length PDGFR and VEGFR have been generated. For cell-based selectivity assays, PAE cells are transfected with fulllength human PDGFR-a, PDGFR-h, or VEGFR-2. Cells are seeded at 4×10⁵ cells/mL in 50 µL growth medium (Ham's F-12 media supplemented with 10% fetal bovine serum, 50,000 units each penicillin and streptomycin, and 500 µg/mL gentamicin) per well in 96-well plates. After 6 to 8 hours, the growth medium is replaced with 50 µL serum-depleted medium (as above, but with 0.1% fetal bovine serum) and cells are incubated overnight. Immediately before compound addition, the medium was replaced with 95 µL serum-depleted medium. Compounds are diluted in 100% DMSO, added to the cells at a final DMSO concentration of 0.25% v/v, and incubated at 37°C for 10 minutes. Cells are stimulated with the appropriate ligand and incubated as above for an additional 8 minutes. The medium is removed and the cells washed once with PBS, then lysed with 50 µL HNTG buffer [20 mmol/L HEPES (pH 7.5), 150 mmol/L NaCl, 2% Triton X-100, 10% glycerol, 5 µmol/L EDTA, 2 mmol/L NaVO₄, and 1 EDTA-free complete protease inhibitor tablet per 25 mL] for 5 minutes at room temperature. Lysates are then diluted with 50 µL HG buffer [20 mmol/L HEPES (pH 7.5), 10% glycerol]. The diluted cell lysates are mixed thoroughly, 50 µL of supernatant are transferred to the ELISA capture plate, and incubated at room temperature for 2 hours with agitation. ELISA capture plates are prepared by coating 96-well ReactiBind goat-antirabbit plates with 100 µL/well of 5 µg/mL rabbit anti-human PDGFR-h, anti-PDGFR-a, or anti-VEGFR-2 antibody for 60 to 90 minutes. At the end of the 2-hour incubation the plates are washed (PBS, 0.1% Tween 20) before incubation with anti-phosphotyrosine-horseradish peroxidase antibody (diluted in PBS, 0.05% Tween 20) for 30 minutes at room temperature. The plates are washed again, then incubated with tetramethylbenzidine and evaluated as described above. IC₅₀ values are calculated as percent inhibition of control. (Only for Reference)

Solubility Information

Solubility	Ethanol: 41.8 mg/mL (100 mM), DMSO: 41.8 mg/mL (100 mM), < 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.3952 mL	11.976 mL	23.9521 mL
5 mM	0.479 mL	2.3952 mL	4.7904 mL
10 mM	0.2395 mL	1.1976 mL	2.3952 mL
50 mM	0.0479 mL	0.2395 mL	0.479 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Roberts WG, et al. Cancer Res, 2005, 65(3), 957-966.

Yang Y L, Cao L B, He W R, et al. Endocytosis triggers V-ATPase-SYK-mediated priming of cGAS activation and

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481