


SAR-20347

Chemical Properties

CAS No. : 1450881-55-6
Formula: C₂₁H₁₈ClFN₄O₄
Molecular Weight: 444.84
Appearance: no data available
Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	SAR-20347 is an inhibitor of TYK2, JAK1/2/3 (IC ₅₀ : 0.6/23/26/41 nM).
Targets(IC ₅₀)	Tyrosine Kinases,JAK
In vitro	SAR-20347 effectively inhibits IL-12-mediated STAT4 phosphorylation (TYK2-dependent, IC ₅₀ : 126 nM) when NK-92 cells are stimulated with IL-12. In the culture media, cells without IL-12 have no measurable IFN-γ, while cells incubated with IL-12 and SAR-20347 demonstrate dose-dependent inhibition of IFN-γ production. SAR-20347 (maximum effect: 5 μM) dose-dependently inhibits the production of secreted embryonic alkaline phosphatase (SEAP).
In vivo	Compared to vehicle-treated animals, SAR-20347 (60 mg/kg) inhibits the production of IFN-γ in the serum by 91%, demonstrating that SAR-20347 can inhibit TYK2 signaling in vivo. SAR-20347 markedly reduces IL-17 production as measured by average signal intensity, consistent with the gene expression analysis.
Kinase Assay	Kinases are prepared in Base Reaction Buffer (20 mM Hepes pH 7.5, 10 mM MgCl ₂ , 1 mM EGTA, 0.02% Brij35, 0.02 mg/mL BSA, 0.1 mM Na ₃ VO ₄ , 2 mM DTT, 1% DMSO) and substrate is added with 1.5 mM CaCl ₂ , 16 μg/mL Calmodulin, and 2 mM MnCl ₂ . Varying concentrations of SAR-20347 in DMSO are added to the kinase reaction along with 10 μM 33P-ATP (activity 0.01 μCi/μL final) for IC ₅₀ determination[1].
Cell Research	Cells are plated in a 96-well v-bottom plate in starvation medium, incubated with SAR-20347 (0.5% DMSO) for 20 minutes at 37°C, 5% CO ₂ , and stimulated with individual cytokines. P-STAT levels are measured in duplicate using MSD plates following the manufacturer's instructions (MSD)[1].
Animal Research	Female 7 to 9-week old C57BL/6 mice are used. Mice are administered vehicle or 50 mg/kg SAR-20347 by oral gavage 30 minutes prior to application of 62.5 mg 5% imiquimod cream or control cream. Another dose of vehicle or 50 mg/kg SAR-20347 is given 5.5 hours following the first dose. This treatment is repeated for 5 days and on day 3 and 4, animals are injected with 100 uL saline to prevent dehydration. On the 6th day, the animals are euthanized and photographs are taken[1].

Solubility Information

A DRUG SCREENING EXPERT

Solubility	DMSO: 100 mg/mL, (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.248 mL	11.240 mL	22.480 mL
5 mM	0.4496 mL	2.248 mL	4.496 mL
10 mM	0.2248 mL	1.124 mL	2.248 mL
50 mM	0.045 mL	0.2248 mL	0.4496 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Kang X, Yang M, Shi Y, et al. Interleukin-15 facilitates muscle regeneration through modulation of fibro/adipogenic progenitors. Cell Communication and Signaling. 2018 Jul 20;16(1):42

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481